

Middlesex University Research Repository

An open access repository of
Middlesex University research

<http://eprints.mdx.ac.uk>

Voermans, A., Mewes, J., van Kaam, Anton, Bayford, Richard ORCID logo ORCID:
<https://orcid.org/0000-0001-8863-6385> and Lepage-Nefkens, I. (2019) PMU31 early
cost-effectiveness analysis of continuous monitoring of lung-aeration with electrical impedance
tomography in preterm neonates with respiratory distress syndrome. Value in Health, Volume
22, Supplement 3, Pages A1-A2, S403-S940. In: ISPOR Europe 2019, 01 Nov 2019,
Copenhagen, Denmark. . ISSN 1098-3015 [Conference or Workshop Item]
(doi:10.1016/j.jval.2019.09.1650)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/28959/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

EARLY COST-EFFECTIVENESS ANALYSIS OF CONTINUOUS MONITORING OF LUNG-AERATION WITH ELECTRICAL IMPEDANCE TOMOGRAPHY IN PRETERM NEONATES WITH RESPIRATORY DISTRESS SYNDROME

Voermans A¹; Mewes J¹; Lepage-Nefkens I¹

1) Panaxea B.V., Amsterdam, the Netherlands

OBJECTIVES: Respiratory distress syndrome (RDS) is relatively common in preterm neonates due to lung immaturity. Clinical management by respiratory support is associated with high complications rates. Guidance on appropriate lung-aeration is limited using conventional thorax X-ray monitoring. Electrical impedance tomography (EIT) allows radiation-free, continuous lung-aeration monitoring to guide effective respiratory support. Clinicians expect EIT implementation to reduce the number of patients requiring mechanical ventilation, and overall complication rates and hospitalisation length. We conducted an early cost-effectiveness analysis of EIT monitoring in preterm neonates with RDS versus standard care in the Netherlands.

METHODS: A decision-analytic model was constructed comparing costs and effects of conventional X-ray versus EIT monitoring for preterm neonates with RDS from the healthcare perspective with a time horizon of two years. Input parameters were based on literature and cost databases. The effects of EIT monitoring were based on consensus by 6 clinical experts for two scenarios, (1) a conservative scenario assuming only a decrease of patients on mechanical ventilation under EIT monitoring, and (2) an optimistic scenario including scenario (1) and assuming an additional 10% relative complication rate decrease in comparison to standard care. Main outcomes were total average costs per patient, number of patients with

bronchopulmonary dysplasia (BPD), and mortality. One-way sensitivity analyses were conducted.

RESULTS: EIT monitoring was estimated to be cost-saving in both scenarios, mainly due to a shorter average hospital length of stay. Total incremental costs per patient for EIT monitored care versus standard care were -€929 and -€10,706 for scenario (1) and (2), respectively. The number of patients with BPD and deaths were reduced. Results were robust to changes in input parameters.

CONCLUSIONS: EIT lung-aeration monitoring in preterm neonates is expected to result in cost-savings and lower mortality and BPD rates, in comparison to standard care, in a Dutch hospital setting.